

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

Uses

- temporarily relieves minor aches and pains due to:
- backache
- headache
- menstrual cramps
- minor pain of arthritis
- muscular aches
- the common cold
- toothache
- temporarily reduces fever

Warnings

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if you have

Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- asthma
- taken a diuretic
- reached age 60 or older

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking aspirin to prevent heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

When using this product

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor.
- children under 12 years: ask a doctor

Other information

Other information

- **each capsule contains:** potassium 20 mg
 - read all warnings and directions before use
 - store at 20 to 25°C (68 to 77°F)
 - avoid excessive heat above 40°C (104°F)
- Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Manufactured for:

Marksans Pharma Inc.

Lake Grove, NY 11755, USA

Manufactured by:

Marksans Pharma Ltd.

Verna, Goa - 403 722, India

Mfg. Lic. No.: GO/Drugs/515

Inactive ingredients

Inactive ingredients Ammonium Hydroxide, FD&C Green No. 3, Gelatin, Iron oxide black, Medium chain triglycerides, Polyethylene glycol, Propylene glycol, Potassium hydroxide, Purified water, Shellac, Sorbitol, Sorbitan monooleate.